

Dedicated to V. I. Arnol'd's 65th birthday

Geometry of the Triangle Equation on Two-Manifolds*

I. A. Dynnikov[†], S. P. Novikov[‡]

Abstract

A non-traditional approach to the discretization of differential-geometrical connections was suggested by the authors in 1997. At the same time we started studying first order difference “black and white triangle operators (equations)” on triangulated surfaces with a black and white coloring of triangles. In this work, we develop the theory of these operators and equations, showing their similarity with the complex derivatives ∂ and $\bar{\partial}$.

Introduction

Do there exist any natural difference analogs of the operators ∂ and $\bar{\partial}$ on the complex plane? No theory of difference “first order” operators that have some properties similar to those of ∂ and $\bar{\partial}$ has been developed before; the attention of the literature is paid only to discrete analogs of the Laplace–Beltrami operator, which are “second order” operators. However, in papers [1, 2, 3] we already studied some “first order” difference operators, starting

*The work of I. Dynnikov is partially supported by Russian Foundation for Fundamental Research, grant no. 02-01-00659, and by Leading Scientific School Support grant no. 00-15-96011; the work of S. Novikov is supported by NSF (USA), grant no. DMS-00772700. Author's contributions are equal; author's names are listed in the alphabetical order.

[†]Mech. and Math. Dept., Moscow State University, Moscow 119992 GSP-2, Russia, e-mail: dynnikov@mech.math.msu.su

[‡]Institute for Physical Sciences and Technology, Univ. of Maryland at College Park, MD 20742, USA; L. D. Landau Institute for Theoretical Physics, Kosygina str. 2, Moscow 117940, Russia, e-mail: novikov@ipst.umd.edu

from the problem of finding a discrete analog of the Laplace transformation for a standard two-dimensional second order linear differential equation. Our operators were originally defined on the equilateral triangular lattice in the space $\mathbb{R}^2$ (see [3]). In [1, 2] these definitions were extended to arbitrary “black and white” triangulations of two-dimensional surfaces without boundary. The “black (white) triangle” equations, which have naturally appeared in the theory, are studied here with respect to their analogy with the operators ∂ and $\bar{\partial}$. It turns out that their properties partially imitate some properties of complex derivatives on two-dimensional manifolds.

1 First order triangle operators on triangulated surfaces. Discrete connections

Let M be a two-dimensional manifold without boundary endowed with a triangulation $\mathcal{T}$. Let $\mathcal{K}$ be a family of triangles from $\mathcal{T}$ such that, for any vertex P of $\mathcal{T}$, there is at least one triangle $T \in \mathcal{K}$ adjacent to P . Let us fix some numeric coefficients $b_{T,P}$ defined for any triangle $T \in \mathcal{K}$ and its vertex $P \in T$.

Definition 1. By a *first order (or triangle) operator* on M associated with the family $\mathcal{K}$ and coefficients $b_{T,P}$ we call an operator $Q^{\mathcal{K}}$ defined by the formula

$$(Q^{\mathcal{K}}\psi)_T = \sum_{P \in T} b_{T,P} \psi_P. \quad (1)$$

The operator $Q^{\mathcal{K}}$ is well-defined as a linear map of the space of all functions ψ_P of a vertex P into the space of functions $\varphi_T = (Q^{\mathcal{K}}\psi)_T$ of a triangle T from the selected family $\mathcal{K}$.

Remark 1. In 1997 (see [1]) we considered operators on multi-dimensional manifolds as well, but in this paper we focus only on the two-dimensional case, where an analogy with the operators ∂ , $\bar{\partial}$ takes place.

In Appendix 2 we provide a generalization of the constructions of this section to higher dimensions.

The maximal family: discrete connections. In the case of the maximal family $\mathcal{K}$ equal to the set of all triangles of the triangulation $\mathcal{T}$, the theory of the equation

$$Q^{\mathcal{K}}\psi = 0 \quad (2)$$


naturally leads to the notions of a *discrete connection* and *curvature*. The coefficients $b_{T,P}$ play the role of the connection coefficients. Following [1], we now introduce the *local holonomy* (or the *curvature*) of the connection $\{b_{T,P}\}$ at a vertex P , which is an operator $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by a matrix of the form

$$K_P = \begin{pmatrix} 1 & k'_P \\ 0 & k''_P \end{pmatrix}. \quad (3)$$

It is constructed as follows.

Let us enumerate all the vertices and triangles included in the star of a vertex P as shown in Fig. 1. We will be constructing a local solution of

Figure 1:



equation (2) in the star of P , starting from arbitrarily chosen values of ψ at vertices P, P_1 . We will have

$$\begin{aligned} \psi_{P_2} &= -\frac{1}{b_{T_1,P_2}}(b_{T_1,P}\psi_P + b_{T_1,P_1}\psi_{P_1}), \\ \psi_{P_3} &= -\frac{1}{b_{T_2,P_3}}(b_{T_2,P}\psi_P + b_{T_2,P_2}\psi_{P_2}), \\ &\dots \\ \tilde{\psi}_{P_1} &= -\frac{1}{b_{T_n,P_1}}(b_{T_n,P}\psi_P + b_{T_n,P_n}\psi_{P_n}) = k'_P\psi_P + k''_P\psi_{P_1}, \end{aligned}$$

where

$$\begin{aligned} k'_P &= \sum_{k=0}^{n-1} (-1)^{k+1} \frac{b_{T_n,P} \cdots b_{T_{n-k},P}}{b_{T_n,P_1} \cdots b_{T_{n-k},P_{n-k+1}}} \\ k''_P &= (-1)^n \frac{b_{T_1,P_1} b_{T_2,P_2} \cdots b_{T_n,P_n}}{b_{T_1,P_2} b_{T_2,P_3} \cdots b_{T_n,P_1}}. \end{aligned}$$

(In the formula for k'_P , we assume $P_{n+1} = P_1$.)

Thus, having started from arbitrary values ψ_P, ψ_{P_1} , after a full turn around the vertex P , we come to new ones

$$(\tilde{\psi}_P, \tilde{\psi}_{P_1}) = (\psi_P, \psi_{P_1}) \begin{pmatrix} 1 & k'_P \\ 0 & k''_P \end{pmatrix}.$$

We say that the connection $\{b_{T,P}\}$ has *zero curvature* at the vertex P if, for any ψ_P, ψ'_P , we have $(\tilde{\psi}_P, \tilde{\psi}_{P_1}) = (\psi_P, \psi_{P_1})$, *i.e.*, if we have


$$k'_P = 0, \quad k''_P = 1.$$

Example 1. Assume that all the connection coefficients are equal to one, $b_{T,P} \equiv 1$, and the number n_P of triangles adjacent to a vertex P is even, $n_P = 2l_P$. Then the holonomy of the connection $\{b_{T,P}\}$ at the vertex P is trivial. Indeed, we have

$$(\psi_P, \psi_{P_{k+1}}) = (\psi_P, \psi_{P_k}) \begin{pmatrix} 1 & -1 \\ 0 & -1 \end{pmatrix}, \quad K_P = \begin{pmatrix} 1 & -1 \\ 0 & -1 \end{pmatrix}^{n_P} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

The *global holonomy* of a discrete connection $\{b_{T,P}\}$ is defined for any “thick” path γ , which is a sequence of triangles $T_1, \dots, T_m$ such that, for all $j = 1, \dots, m-1$, the triangles T_j and T_{j+1} have a common edge $\kappa_j = T_j \cap T_{j+1}$, whose vertices are denoted by P'_j, P''_j (see Fig. 2). This is done as follows.

Figure 2: A thick path



Start from a solution $(\psi_P, \psi_{P'_1}, \psi_{P''_1})$ of equation (2) in the triangle T_1 :

$$b_{T_1,P}\psi_P + b_{T_1,P'_1}\psi_{P'_1} + b_{T_1,P''_1}\psi_{P''_1} = 0.$$

We assume that ψ_P and $\psi_{P'_1}$ are chosen arbitrarily, and $\psi_{P''_1}$ is found from the equation. Then we extend the solution along γ successively to the triangles

$T_2, T_3, \dots$, *i.e.*, to the vertices $P'_2, P''_2, P'_3, P''_3, \dots$. This can be done in a unique way. If γ is a loop, *i.e.*, $T_{m+1} = T_1$, then, having passed over γ , we will obtain a new solution $(\tilde{\psi}_P, \tilde{\psi}_{P'_1}, \tilde{\psi}_{P''_1})$ of (2) in the triangle T_1 . Thus, we have constructed the *holonomy operator* $R_\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$:

$$(\psi_P, \psi_{P'_1}) \mapsto (\tilde{\psi}_P, \tilde{\psi}_{P'_1}) = (\psi_P, \psi_{P'_1}) \cdot R_\gamma.$$

Lemma 1. *The global holonomy of any discrete connection $\{b_{P,T}\}$ with zero curvature is a well defined homomorphism $\pi_1(M, T_1) \rightarrow GL(2, \mathbb{R})$.*

Proof. For any two adjacent triangles T, T' , *i.e.*, ones having exactly one common edge, we denote by $R_{T,T'}$ the extension operator of a solution of (2) from the triangle T to T' : $\psi^{(T)} \mapsto \psi^{(T')} = \psi^{(T)} \cdot R_{T,T'}$. By definition, we have

$$R_\gamma = R_{T_1, T_2} \cdot \dots \cdot R_{T_{m-1}, T_m} \cdot R_{T_m, T_{m+1}}.$$

Clearly, the correspondence $\gamma \mapsto R_\gamma$ is multiplicative:

$$R_{\gamma_1 \circ \gamma_2} = R_{\gamma_1} \circ R_{\gamma_2},$$

and consistent with the inversion:

$$R_{\gamma^{-1}} = (R_\gamma)^{-1}.$$

We have only to check the invariance of R_γ under a homotopy of the path γ .

Any homotopy of γ can be reduced to finitely many insertions and removals of fragments like this:

$$\begin{aligned} \dots, T, T', T, \dots &\leftrightarrow \dots, T, \dots, \\ \dots, T, T', T'', \dots, T^{(n)}, T, \dots &\leftrightarrow \dots, T, \dots, \end{aligned}$$

where $T, T', \dots, T^{(n)}$ are triangles from the star of a vertex listed in the order we meet them when walking around the vertex. The holonomy operator R_γ is unchanged under the first operation in view of $R_{T,T'} R_{T',T} = \text{id}$, and it does so under the latter in view of the triviality of the local holonomy. $\square$

Theorem 1. *For any triangulation $\mathcal{T}$ of a surface M and any homomorphism $\rho : \pi_1(M) \rightarrow GL(2, \mathbb{R})$ there exists a zero curvature discrete connection $\{b_{T,P}\}$ that generates the representation ρ .*

See Appendix 1 for a proof of this theorem.


In the sequel, we will consider in detail only one special case in which the star of each vertex of $\mathcal{T}$ consists of an even number of triangles and all the connection coefficients are equal to 1, $b_{T,P} \equiv 1$. As we have already seen, in this case, the connection $\{b_{T,P}\}$ has zero curvature, and hence, defines a homomorphism $\rho : \pi_1(M) \rightarrow GL(2, \mathbb{R})$. We call this connection *canonical*.

Proposition 1. *The holonomy group of the canonical connection associated with a triangulation all whose vertices have an even valence is a subgroup of the group $S_3 \subset GL(2, \mathbb{R})$ generated by matrices*

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \begin{pmatrix} -1 & 0 \\ -1 & 1 \end{pmatrix}. \quad (4)$$

Proof. Let γ be a thick path $T_1, T_2, \dots, T_{m+1} = T_1$. Let us color the vertices of T_1 into three distinct colors, say, a, b, c , and then continue the coloring so that all three vertices of any triangle T_j , $j = 2, 3, \dots$, have different colors (see Fig. 3). At the same time, we extend a solution ψ from the triangle

Figure 3: Coloring vertices in three colors



T_1 to $T_2, T_3, \dots$. Since the sum of the values of ψ over the vertices of any triangle must be zero, we will have $\psi_P = \psi_{P'}$ for any two vertices P, P' of the same color. We denote by ψ_a, ψ_b, ψ_c the values of ψ at vertices of the corresponding color. We have $\psi_a + \psi_b + \psi_c = 0$.

After passing over γ , we come to another coloring of the vertices of T_1 , which is obtained from the original one by a permutation from S_3 . The group S_3 is generated by the permutations $\begin{pmatrix} a & b & c \\ b & a & c \end{pmatrix}$ and $\begin{pmatrix} a & b & c \\ c & b & a \end{pmatrix}$. For the first permutation, we have

$$(\tilde{\psi}_a, \tilde{\psi}_b) = (\psi_b, \psi_a) = (\psi_a, \psi_b) \cdot \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

for the latter one, we have

$$(\tilde{\psi}_a, \tilde{\psi}_b) = (-\psi_a - \psi_b, \psi_b) = (\psi_a, \psi_b) \cdot \begin{pmatrix} -1 & 0 \\ -1 & 1 \end{pmatrix}.$$

□

In the case of $\{b_{T,P}\}$ the canonical connection, we call equation (2) the *triangle equation*:

$$(Q\psi)_T = \sum_{P \in T} \psi_P = 0. \quad (5)$$

In order to construct a solution, we have to take initial values ψ_a, ψ_b invariant under all holonomy operators. According to Proposition 1, there are three cases:

Case 1 : the holonomy group G is isomorphic either to S_3 or S_3^+ (the group of even permutations); the representation $G \rightarrow GL(2, \mathbb{R})$ is irreducible;

Case 2 : the holonomy group is isomorphic to $\mathbb{Z}_2$; up to multiplicative constant, there is exactly one invariant vector;

Case 3 : the holonomy group is trivial; all vectors from $\mathbb{R}^2$ are invariant.

Thus, we have proved the following.

Theorem 2. *The space of “covariant constants”, i.e., of solutions of the triangle equation, is:*

two-dimensional if the holonomy group is trivial;

one-dimensional if the holonomy group is isomorphic to $\mathbb{Z}_2$;

trivial in the other cases.

Now consider the operator $L = Q^+Q$, where Q is the first order operator associated with the canonical connection on the surface M . The operator L can be written as follows:

$$L = -2\delta d + 3n_P = -2\Delta + 3n_P, \quad (6)$$

where n_P is the “potential” equal to the valence of a vertex, d, δ are the boundary and coboundary operators, respectively, $d = \delta^+$.

Proposition 2. *The operator L has a non-trivial space of zero modes $L\psi = 0$ if and only if the holonomy group of the canonical connection is either trivial, in which case the space of zero modes is two-dimensional, or isomorphic to $\mathbb{Z}_2$, in which case it is one-dimensional. These modes ψ satisfy the equation $Q\psi = 0$.*

Remark 2. For $\mathcal{T}$ a uniform triangulation, i.e., such that all vertices have the same valence $n_P = 2l = \text{const}$, the zero modes of L are eigenfunctions of the Laplace–Beltrami operator Δ with maximal eigenvalue.

It is interesting to compare our constructions with some results previously obtained in graph theory (see [4]). Let us consider the Poincaré dual graph Γ of our triangulation. Vertices of Γ correspond to triangles of $\mathcal{T}$. This graph Γ is “dichromatic” in the graph-theoretical terminology if and only if there exists a coloring of triangles of $\mathcal{T}$ in black and white such that any two adjacent triangles have different colors. The Laplace–Beltrami operators on dichromatic graphs Γ were studied in [4]. The space $\mathcal{L}_\Gamma$ of functions depending on vertices of Γ splits into the direct sum

$$\mathcal{L}_\Gamma = \mathcal{L}_b \oplus \mathcal{L}_w,$$

where $\mathcal{L}_b$ (respectively, $\mathcal{L}_w$) is the space of functions of black (respectively, white) triangles of $\mathcal{T}$.

In [1] we studied operators $Q_{bw} : \mathcal{L}_w \rightarrow \mathcal{L}_b$, and the operators $Q_{wb} = Q_{bw}^+$ adjoint to them, associated with the coloring. They were needed for a generalization of the Laplace transformation. This was a new type of first order difference operators, except in the following case: for an equilateral triangular lattice with the natural coloring, both spaces $\mathcal{L}_w, \mathcal{L}_b$ are canonically isomorphic to the space of functions depending on vertices (see below). In our previous work [1], “second order Schrödinger operators” of the form $L = Q_{wb}Q_{wb}^+$ and $L' = Q_{wb}^+Q_{wb}$ acting on the spaces $\mathcal{L}_w$ and $\mathcal{L}_b$, respectively, were discussed.

It is easy to see that the standard graph-theoretical Laplace–Beltrami operator on the dual graph Γ has a matrix of the form:

$$\Delta_\Gamma = \begin{pmatrix} 0 & Q_{wb} \\ Q_{wb}^+ & 0 \end{pmatrix}, \quad \Delta_\Gamma^2 = \begin{pmatrix} Q_{wb}Q_{wb}^+ & 0 \\ 0 & Q_{wb}^+Q_{wb} \end{pmatrix} = L \oplus L'.$$

So the results obtained in the work of P. Sarnak [4] involve, in fact, the operators of the type Q_{wb} but not of the type studied in the present work.

Let us ask the following

Question: when a triangulation $\mathcal{T}$ of a surface M admits a black and white (or b/w) coloring of triangles such that any two adjacent triangles are of different colors?

The answer, which is obvious, is given in the following Lemma.

Lemma 2. *A b/w coloring of a triangulation $\mathcal{T}$ exists if and only if any thick loop $\gamma = (T_1, \dots, T_m, T_{m+1} = T_1)$, where $T_i \cap T_{i+1}$ is an edge for all $i = 1, \dots, m$, has an even length: $m = 2k$.*

Thus, we have the following three homomorphisms $\rho_1, \rho_2, \rho_3 : \pi_1(M) \rightarrow \mathbb{Z}_2$:

- 1) the parity ρ_1 of the global holonomy, $\pi_1(M) \rightarrow S_3 \rightarrow \mathbb{Z}_2$;
- 2) the orientation homomorphism ρ_2 ;
- 3) the parity homomorphism ρ_3 of the number of triangles in a thick loop.

Lemma 3. *We have $\rho_1 \rho_2 = \rho_3$.*

We skip the easy proof.

Corollary 1. *a) Let M be orientable; then a b/w coloring of $\mathcal{T}$ exists if and only if the holonomy group of the canonical connection associated with $\mathcal{T}$ is either trivial or isomorphic to the group S_3^+ of even permutations;*

b) Let M be non-orientable; then a b/w coloring exists if and only if the holonomy group is isomorphic to $\mathbb{Z}_2$ and we have $\rho_1 = \rho_2$.

In the case $\rho_3 = 1$, we fix a b/w coloring of $\mathcal{T}$ and consider the first order operators Q_b and Q_w of the form (1) associated with the families of black and white triangles, respectively. The corresponding equations $Q_b \psi = 0$ and $Q_w \psi = 0$ are called *black triangle equation* and *white triangle equation*, respectively. Clearly, we have

$$Q_b^+ Q_b = -\delta d + \frac{3}{2} n_P = Q_w^+ Q_w. \quad (7)$$

2 Maximum principle

As before, let M be a surface endowed with a triangulation $\mathcal{T}$. Let $M' \subset M$ be a surface (with boundary) that has a form of a finite simplicial subcomplex

of M . Assume that M' is orientable and the holonomy of the canonical connection of M' is globally trivial.

As we saw in the previous Section, there exists a b/w coloring of triangles from M' such that any two adjacent triangles have different colors. Also, there exists a coloring of vertices into three colors, a, b, c , such that all three vertices of any triangle are of different colors. The space V of covariant constants is two-dimensional and consists of functions ψ_P that depend only on the color of a vertex P , and the values ψ_a, ψ_b, ψ_c of ψ at vertices of the corresponding color satisfy the relation $\psi_a + \psi_b + \psi_c = 0$.

Let $\mathcal{K}$ be the set of black triangles in M' . For any solution ψ of the black triangle equation $Q^{\mathcal{K}}\psi = 0$, we define a mapping $\widehat{\psi} : \mathcal{K} \rightarrow V \cong \mathbb{R}^2$ from the set of black triangles to the space of covariant constants on M' as follows. For a black triangle $T \subset M'$, we define $\widehat{\psi}(T)$ to be the covariant constant that coincides with ψ on vertices of T . In other words, $\widehat{\psi}(T) = (\psi_{P_a(T)}, \psi_{P_b(T)}, \psi_{P_c(T)})$, where $P_a(T), P_b(T), P_c(T)$ are vertices of T of colors a, b, c , respectively.

Let us denote by $\partial_- M'$ the set of triangles in M' that are adjacent to the boundary $\partial M'$, *i.e.*, such that $T \subset M'$ and $T \cap \partial M' \neq \emptyset$.

Theorem 3 (Maximum principle). *For any solution ψ of the black triangle equation on a submanifold $M' \subset M$, the image $\widehat{\psi}(\mathcal{K})$ of the set of black triangles in the space of covariant constants is contained in the convex hull of the image of the boundary $\widehat{\psi}(\partial_- M')$.*


Proof. We assume that ψ is not a covariant constant, in which case the assertion is trivial because the image $\widehat{\psi}(\mathcal{K})$ is just one point $\psi \in V$.

Since the complex M' is finite, the convex hull C of $\widehat{\psi}(\mathcal{K})$ is a polygon in the plane $V \cong \mathbb{R}^2$. The assertion means that the corners of C are the images of boundary triangles $T \in \mathcal{K} \cap (\partial_- M')$. In order to prove this, we are going to show that the image of any internal black triangle, *i.e.*, a triangle $T \in \mathcal{K} \setminus (\partial_- M')$, is not a corner of C .

For an internal black triangle $T \subset M'$, let us denote by $T_1, \dots, T_6$ the six black triangles that are at distance 2 from T in the sense of thick paths and are arranged as shown in Fig. 4. (The pairs of triangles (T_1, T_6) , (T_2, T_3) , (T_4, T_5) may coincide, which will not affect the proof.) If T is an internal triangle of M' , then all six triangles $T_1, \dots, T_6$ lie in M' . Without loss of generality we may assume that vertices are colored as showing in Fig. 4.


There are three families of parallel straight lines in the space of covariant constants: $\psi_a = \text{const}$, $\psi_b = \text{const}$, and $\psi_c = \text{const}$. If two black triangles

Figure 4:



$T, T' \subset M'$ have a common vertex of color $x \in \{a, b, c\}$, then the points $\widehat{\psi}(T), \widehat{\psi}(T') \in V$ lie in a straight line $\psi_x = \text{const}$. In Fig. 5, one of possible arrangements of points $\widehat{\psi}(T), \widehat{\psi}(T_1), \dots, \widehat{\psi}(T_6)$, which are marked simply as 0, 1, 2, 3, 4, 5, 6, respectively, is shown. It is easy to see that, for any other

Figure 5:



possible arrangement, the point $\widehat{\psi}(T)$ lies between either $\widehat{\psi}(T_1), \widehat{\psi}(T_6)$, or $\widehat{\psi}(T_2), \widehat{\psi}(T_3)$, or $\widehat{\psi}(T_4), \widehat{\psi}(T_5)$. $\square$


In the particular case $M' = M$, we come to the following

Corollary 2. *Let M be a closed triangulated surface such that the holonomy group of the canonical connection is trivial. Then the only solutions of the black (or white) triangle equation on M are covariant constants, which form a two-dimensional space.*

3 Triangle equations in the Euclidean geometry

From now on, we will consider the case when our surface M is the Euclidean plane $\mathbb{R}^2$ triangulated so that all triangles are equilateral. Its vertices form a two-dimensional lattice $\ell \subset \mathbb{R}^2$, $\ell \cong \mathbb{Z}^2$. We denote by t_1 and t_2 the basis shifts of the lattice. Each triangle of this triangulation is either of the form $\langle P, t_1 \cdot P, t_2 \cdot P \rangle$ (in which case we call it *white* and denote by T_P^w), or of the form $\langle P, t_1^{-1} \cdot P, t_2^{-1} \cdot P \rangle$ (in which case we call it *black* and denote by T_P^b), see Fig. 6.

Figure 6: Black and white coloring of the Euclidean plane



Thus, in this case, there is a natural one-to-one correspondence between vertices of the triangulation and black (white) triangles: $P \leftrightarrow T_P^b$ (respectively, $P \leftrightarrow T_P^w$). Due to this correspondence, we can think of operators Q_b, Q_w of the form (1) associated with the set of black or white triangles, respectively, as ones mapping the space of functions on the lattice $\mathbb{Z}^2$ to itself. In other words, Q_b, Q_w have the form

$$\begin{aligned} Q_w &= x_n + y_n t_1 + z_n t_2, \\ Q_b &= u_n + v_n t_1^{-1} + w_n t_2^{-1}, \end{aligned} \tag{8}$$

where $n = (n_1, n_2) \in \mathbb{Z}^2$. Notice, that operators of “black” type Q_b are (formally) conjugated to operators of “white” type Q_w , as we have $t_j^+ = t_j^{-1}$.

Consider a real self-adjoint second order (Schrödinger) operator

$$(L\psi)_P = \sum_{P'} b_{P,P'} \psi_{P'}, \quad b_{P,P'} = b_{P',P}, \tag{9}$$

where $b_{P,P'} > 0$ holds if either we have $P = P'$ or $\langle P, P' \rangle$ is an edge, and we have $b_{P,P'} = 0$ otherwise. In other words, L is an operator of the form

$$L = a_n + b_n t_1 + c_n t_2 + d_n t_1^{-1} t_2 + e_n t_1^{-1} + f_n t_2^{-1} + g_n t_1 t_2^{-1}. \quad (10)$$

It admits a unique factorization of the form

$$L = Q_b^+ Q_b + U_P \quad (11)$$

and of the form

$$L = Q_w^+ Q_w + V_P, \quad (12)$$

where all coefficients of Q_b and Q_w are positive.

Interesting classes of such operators and their spectral theory in the space $\mathcal{L}_2(\mathbb{Z}^2)$ are studied in [1, 3]. In particular, the attention is paid to zero modes of the operator $Q_b^+ Q_b$ (respectively, $Q_w^+ Q_w$), which are functions $\psi \in \mathcal{L}_2(\mathbb{Z}^2)$ such that $Q_b \psi = 0$ (respectively, $Q_w \psi = 0$). Explicit solutions are found in the case of coefficients of the form $\exp(l_j(n))$, where $l_j(n) = l_{j1}n_1 + l_{j2}n_2$, $j = 1, 2$, are linear forms:

$$Q_w = 1 + ce^{l_1(n)}t_1 + de^{l_2(n)}t_2 = Q(c, d). \quad (13)$$

Especially interesting is the case when $l_{ij} + l_{ji} = h$ is independent of i, j . In this case, the operators $Q(c, d)$ satisfy the relation

$$Q(c, d)^+ Q(c, d) - 1 = q^2 (Q(c/q^2, d/q^2) Q(c/q^2, d/q^2)^+ - 1), \quad (14)$$

where $q = e^{l_{11}} = e^{l_{22}} = e^{(l_{12} + l_{21})/2}$. In some cases, these relations allow to establish a wonderful property of the spectrum of operators $L = Q^+ Q$ and $L' = Q Q^+$ in the Hilbert space $\mathcal{L}_2(\mathbb{Z}^2)$ similar to that of the famous ‘‘Landau operator’’ in the magnetic field, though there is nothing like a physical magnetic field in our situation. This phenomena are coming only from the nature of the discrete systems.

A pair of operators (Q_b, Q_w) defines a discrete connection on the plane $\mathbb{R}^2$ as described in Section 1. As shown in [1], this connection has trivial local holonomy in the special case $l_{ij} + l_{ji} = 0$.

It is also noticed in [1] that operators $Q_w = 1 + y_n t_1 + z_n t_2$ and $Q_b = 1 + v_n t_1^{-1} + w_n t_n^{-1}$ define a connection of zero curvature if and only if there exists an everywhere non-zero function f such that

$$((Q_w - 1)(Q_b - 1) - 1) = f \cdot ((Q_b - 1)(Q_w - 1) - 1). \quad (15)$$

4 Discrete analog of the operators ∂ and $\bar{\partial}$

Recall that we consider an equilateral triangular lattice $\ell \cong \mathbb{Z}^2$ in the Euclidean plane $\mathbb{R}^2$ and the corresponding b/w triangulation: the triangles $T_n^w = \langle (n_1, n_2), (n_1 + 1, n_2), (n_1, n_2 + 1) \rangle$ are white, and the triangles $T_n^b = \langle (n_1, n_2), (n_1 - 1, n_2), (n_1, n_2 - 1) \rangle$ are black, where $n = (n_1, n_2) \in \mathbb{Z}^2$.

We will consider the following operators on the space of functions on ℓ :

$$Q = 1 + t_1 + t_2, \quad Q^+ = 1 + t_1^{-1} + t_2^{-1}, \quad (16)$$

which are formally conjugate to each other, and show that they have many properties similar in a sense to those of the operators $\partial = \partial/\partial z$ and $\bar{\partial} = \partial/\partial \bar{z}$ on the complex plane.

We denote by H the kernel of the operator Q^+ . It will play a role of the space of *holomorphic functions*. It consists of functions ψ on ℓ that have zero sum over all the vertices of any black triangle: $Q^+\psi = 0$. We have not found any natural algebra structure on H , but we have constructed analogs of polynomials, the Taylor expansion, and the Cauchy formula. A discrete version of the maximum principle, which was considered in Section 2 in a more general situation, also takes place.

Now we introduce the space $\mathcal{P}_k$ of “polynomials of degree $\leq k$ ”, which are solutions of the following system of equations:

$$Q^+\psi = 0, \quad Q^{k+1}\psi = 0. \quad (17)$$

Proposition 3. *The dimension of the subspace $\mathcal{P}_k$ equals $2k + 2$.*

Proof. By definition, $\mathcal{P}_0$ is the space of functions ψ satisfying both equations

$$Q\psi = 0, \quad Q^+\psi = 0. \quad (18)$$

It means that ψ is a covariant constant with respect to the canonical connection of our triangulation. As we have seen in Section 1, in the case of trivial holonomy, the space of covariant constants is two-dimensional. Thus, we have

$$\dim \mathcal{P}_0 = 2.$$

Notice that, since the operators Q and Q^+ commute, the space H of “holomorphic” functions is invariant under Q : $Q(H) \subset H$. We claim that

we also have $Q(H) \supset H$, i.e., for any function $\varphi \in H$, the following system of equations is consistent:

$$Q\psi = \varphi, \quad Q^+\psi = 0. \quad (19)$$

Indeed, the system defines an *affine* discrete connection on the plane similar to the canonical connection from Section 1. One can easily check that the local holonomy of this connection is trivial if and only if we have $Q^+\varphi = 0$, which holds by assumption. The rest of the argument is the same as in the case of linear discrete connection.

Thus, the restriction of Q to the space H is a linear operator whose image is the whole space H and the kernel is two-dimensional. It follows immediately that

$$\dim(\mathcal{P}_{k+1}/\mathcal{P}_k) = 2.$$

□

Remark 3. One can show that, for any $\psi \in \mathcal{P}_k \setminus \mathcal{P}_{k-1}$, there exists a degree $k-1$ polynomial $u_{k-1}(z, \bar{z})$ with complex coefficients such that the following holds

$$\psi_n = \operatorname{Re} \left(\alpha e^{\frac{2\pi i}{3}(n_1 - n_2)} (z^k + u_{k-1}(z, \bar{z})) \right), \quad (20)$$

where $n = (n_1, n_2)$, $z = n_1 + e^{\frac{2\pi i}{3}} n_2$, $\alpha \in \mathbb{C}$.

Now we introduce an analog of the Taylor expansion for functions from H . We will need to consider “big black triangles” $T_n^{(k)} = \langle (n_1, n_2), (n_1 - 2k - 1, n_2), (n_1, n_2 - 2k - 1) \rangle$, where $n = (n_1, n_2) \in \ell$. A big black triangle is homothetic to an ordinary black triangle and has $2k + 2$ vertices of the lattice in each side.

We need the following preparatory lemma.

Lemma 4. *For any “holomorphic function” $\psi \in H$, any $n \in \ell$ and $k \geq 0$, there exists a unique “degree $\leq k$ polynomial” $p_k \in \mathcal{P}_k$ such that p_k coincides with ψ in $T_n^{(k)}$.*

Proof. We prove the existence by induction. For $k = 0$, the assertion is true, since there always exists a covariant constant p_0 coinciding with ψ in just one black triangle.

Assume that the assertion is proved for $k = l - 1$. Then we can find $p_{l-1} \in \mathcal{P}_{l-1}$ such that $p_{l-1} = Q\psi$ wherever in $T_{(n_1-1, n_2-1)}^{(l-1)}$. As we have already seen, we can find a function $\varphi \in H$ such that $Q\varphi = p_{l-1}$, and a covariant

constant $p_0 \in \mathcal{P}_0$ such that $p_0 = \psi - \varphi$ in $T_n^b = T_n^{(0)}$. We set $p_l = \varphi + p_0$. By construction, we will have: $p_l \in \mathcal{P}_l$, $p_l = \psi$ in $T_n^{(0)}$ and $Qp_l = Q\psi$ in $T_{(n_1-1, n_2-1)}^{(l-1)}$.

Thus, the difference $p_l - \psi$ satisfies the following

$$Q^+(p_l - \psi) = 0 \text{ everywhere, } \quad Q(p_l - \psi) = 0 \text{ in } T_{(n_1-1, n_2-1)}^{(l-1)},$$

which means that the sum of the values of the function $p_l - \psi$ over the vertices of any triangle, black or white, contained in $T_n^{(l)}$ is zero. Since $p_l = \psi$ in $T_n^{(0)}$, this implies that the function $p_l - \psi$ is identically zero in $T_n^{(l)}$.

The uniqueness of the “polynomial” p_k follows from the dimension argument: the dimension of $\mathcal{P}_k$ is $2k + 2$ by Proposition 3, and the $2k + 2$ values of ψ at vertices lying in one side of $T_n^{(k)}$, say, points $(n_1 - j, n_2)$, where $j = 0, 1, \dots, 2k + 1$, can be arbitrary as implied by the following lemma. $\square$

Lemma 5. *Let $Y_{n_1 n_2}$ be the following subset of the lattice ℓ :*

$$Y_n = \{(n_1, n_2)\} \cup \{(n_1 - j, n_2), (n_1, n_2 + j), (n_1 + j, n_2 - j)\}_{j \geq 1}, \quad (21)$$

where $n = (n_1, n_2)$. Let V_n be the linear space of all real functions on Y_n . Then, for any $n \in \ell$ the restriction map $H \rightarrow V_n$ is an isomorphism.

In other words, in order to get a solution to the equation $Q^+\psi = 0$ one can set values of ψ in all points from Y_n arbitrarily, and then the equation will prescribe the values of ψ in all the other points.


Proof. The three rays with the origin n at which all the points from Y_n lie, cut the plane into three parts, which are cyclically interchanged under rotation by $2\pi/3$ around n . Consider one of these parts $U = \{(n_1 - j_1, n_2 + j_2)\}_{j_1, j_2 > 0}$. All black triangles in U have the form $T_{(n_1-j_1+1, n_2+j_2)}^b$, where $j_1, j_2 > 0$. The corresponding equations are:

$$\psi_{n_1-j_1, n_2+j_2} = -\psi_{n_1-(j_1-1), n_2+j_2} - \psi_{n_1-(j_1-1), n_2+(j_2-1)}. \quad (22)$$

They allow to express recursively the value of ψ at any point from $\ell \cap U$ via the values of ψ at $\ell \cap \partial U$.

A similar situation takes place in the other two parts of the plane that are obtained from U by rotation by $2\pi/3$ around n . The recursion process is illustrated in Fig. 7, where the vertices from Y_n are marked by bold circles. $\square$

Figure 7: Recursive construction of ψ




There is a canonical way to associate three “polynomials of degree k ” $p_{k,1}, p_{k,2}, p_{k,3}$ with any big black triangle $T_n^{(k)}$ by setting them to be identically zero everywhere in $T_n^{(k)}$ except at vertices lying in one of the sides where they are equal to ± 1 , namely:

$$\begin{aligned} p_{k,1}(n_1 - 2k - 1 + j, n_2) &= (-1)^{j+k}; \\ p_{k,2}(n_1, n_2 - j) &= (-1)^{j+k}; \\ p_{k,3}(n_1 - j, n_2 - 2k - 1 + j) &= (-1)^{j+k}, \end{aligned} \quad (23)$$

$j = 0, 1, \dots, 2k + 1$. In Fig. 8 the values of $p_{k,1}$ in $T_n^{(k)}$ are shown for $k = 1, 2$. The pictures for $p_{k,2}$ and $p_{k,3}$ are obtained by rotation by $2\pi/3$. Obviously,

Figure 8: “Polynomials” $p_{k,1}$ in $T_n^{(k)}$; $k = 1, 2$



we have

$$Qp_{k,i} = p_{k-1,i}, \quad (24)$$

where $p_{k,i}$ is associated with $T_{(n_1,n_2)}^{(k)}$, and $p_{k-1,i}$ with $T_{(n_1-1,n_2-1)}^{(k-1)}$.


The polynomials $p_{k,j}$ are linearly dependent:

$$p_{k,1} + p_{k,2} + p_{k,3} \in \mathcal{P}_{k-1}, \quad (25)$$

where we assume that the polynomials are associated with the same big triangle $T_n^{(k)}$. Thus, in order to get a basis in the space of polynomials we have to select a big triangle $T_n^{(k)}$ for each k and a pair of polynomials $p_{k,i}, p_{k,j}$, $i, j \in \{1, 2, 3\}$. We do it as follows.

For a big triangle $T_{(n_1,n_2)}^{(k)}$, we define its *two-side extension* of type (12), (23), or (31) to be the big triangle $T_{(n_1+1,n_2+1)}^{(k+1)}$, $T_{(n_1+1,n_2)}^{(k+1)}$, or $T_{(n_1,n_2+1)}^{(k+1)}$, respectively (see Fig. 9).

Figure 9: Two-side extensions



Let us fix a sequence of big triangles $T(0), T(1), \dots$ such that

- 1) $T(0)$ is an ordinary black triangle;
- 2) $T(k+1)$ is a two-side extension of $T(k)$ (thus, $T(k) = T_n^{(k)}$ for some $n \in \ell$);
- 3) the union $\bigcup_k T(k)$ is the whole plane $\mathbb{R}^2$.

Such a sequence will be called *admissible*.

With an admissible sequence of triangles we associate a basis ψ_j^1, ψ_j^2 , $j = 0, 1, \dots$, in the space $\bigoplus_k \mathcal{P}_k$ of polynomials in the following way. If $T(k)$ is the two-side extension of $T(k-1)$ of type (ij) , we set ψ_k^1 and ψ_k^2 to be the polynomials $p_{k,i}$ and $p_{k,j}$ associated with the triangle $T(k)$. We also denote by $T^b(k)$ the (ordinary) black triangle $T_{(n_1-k,n_2-k)}^b$, where $T(k) = T_{(n_1,n_2)}^{(k)}$.

By construction, we have $\psi_j^i = 0$ everywhere in $T(k)$ if $k < j$. Thus, any series of the form

$$\sum_{k=0}^{\infty} (\alpha_k^1 \psi_k^1 + \alpha_k^2 \psi_k^2) \quad (26)$$

converges everywhere in ℓ , since, by the definition of admissible sequence, for any $n \in \ell$, we have $n \in T(k)$ for any sufficiently large k .

Theorem 4. *For any admissible sequence of triangles $T(0), T(1), \dots$ and any “holomorphic” function $\psi \in H$, there exists a unique series of coefficients $\alpha_0^1, \alpha_0^2, \alpha_1^1, \alpha_1^2, \alpha_2^1, \alpha_2^2, \dots$ such that the series (26) converges to the function ψ . Moreover, the coefficients α_k^1, α_k^2 can be found from the knowledge of the value of the “ k th derivative” $Q^k\psi$ at the vertices of the triangle $T^b(k)$.*

Proof. The first assertion of the theorem follows from the fact that we can approximate ψ in the triangle $T(k)$ by a “polynomial” of degree k and from $\lim_{k \rightarrow \infty} T(k) = \mathbb{R}^2$.

It is straightforward to check that, if ψ is identically zero in $T(k)$, then $Q^k\psi$ is identically zero in $T^b(k)$. Thus, $Q^j\psi_k^i$ is not identically zero in $T^b(k)$ if and only if $j = k$. This implies the latter assertion of the theorem. $\square$

Now we construct an analog of the Cauchy formula for recovering a solution ψ of the equation $Q^+\psi = 0$ in a domain from the knowledge of ψ at the boundary of the domain.

By a domain we will always mean a closed domain D in the plane such that D is a simplicial subcomplex of our regular triangulation. For a black or white triangle T , we will write $T \in \partial_+ D$ if we have $T \not\subset D$ and $T \cap \partial D \neq \emptyset$.

Consider the following function (the famous Pascal triangle, see Fig. 10).

$$G_{(n_1, n_2)} = \begin{cases} (-1)^{n_1+n_2} \binom{n_1}{n_1+n_2} & \text{if } n_1, n_2 \geq 0, \\ 0 & \text{otherwise.} \end{cases} \quad (27)$$


Lemma 6. *The function G is a fundamental solution to the black triangle equation, i.e., we have*

$$Q^+G = \delta, \quad (28)$$

where δ is the delta-function

$$\delta_n = \begin{cases} 1 & \text{if } n = (0, 0), \\ 0 & \text{otherwise.} \end{cases}$$

Figure 10: The function G



Proof. The formula (27) can be rewritten in the form

$$G_n = \sum_{k=0}^{\infty} ((-t_1^{-1} - t_2^{-1})^k \delta)_n. \quad (29)$$

Applying $Q^+ = (1 + t_1^{-1} + t_2^{-1})$ to both sides of (29) we obtain (28). $\square$

Proposition 4. *Let ψ be a “holomorphic function” in a finite domain D , i.e., a solution of the equation $Q^+\psi = 0$ in D . Then, for any $n \in D \cap \ell$, the following holds*

$$\psi_n = \sum_{T_m^b \in \partial_+ D} (Q^+\psi)_m G_{n-m}, \quad (30)$$

where we set $\psi_m = 0$ for all $m \notin D$.

Note that the right hand side of (30) involves only the values of ψ at the boundary ∂D .

Proof. Notice that, if $T_m^b \notin \partial_+ D$, then we have $(Q^+\psi)_m = 0$. Indeed, if $T_m^b \subset D$, then the equality holds by assumption, and if $T_m^b \cap D = \emptyset$, it holds, since ψ is assumed to be identically zero outside D . So, the right hand side of (30) can be written as

$$\sum_{m \in \ell} (Q^+\psi)_m G_{n-m}, \quad (31)$$

Denote this sum by $\tilde{\psi}_n$.

We may assume without loss of generality that the domain D is contained in the sector $n_1, n_2 > 1$. Then, for any $n = (n_1, n_2)$ such that $T_n^b \in \partial_+ D$, we have $n_1, n_2 > 0$. This implies

$$\tilde{\psi}_n = \psi_n = 0 \quad \text{if } n_1 \leq 0 \text{ or } n_2 \leq 0. \quad (32)$$

From (28) we have

$$\begin{aligned} (Q^+(\psi - \tilde{\psi}))_n &= (Q^+\psi)_n - \sum_{m \in \ell} (Q^+\psi)_m (Q^+G)_{n-m} \\ &= (Q^+\psi)_n - \sum_{m \in \ell} (Q^+\psi)_m \delta_{n-m} \\ &= (Q^+\psi)_n - (Q^+\psi)_n = 0. \end{aligned}$$

Thus, the function $\psi - \tilde{\psi}$ belongs to H and, according to (32), vanishes at Y_0 . It follows from Lemma 5 that $\psi - \tilde{\psi} = 0$. $\square$

In this proof, we have used two properties of the function G : the first one is relation (28), and the second one is that $G = 0$ outside the sector $n_1, n_2 \geq 0$. However, the latter is not needed as will follow from the next lemma.

Lemma 7. *For any function $\psi : \ell \rightarrow \mathbb{R}$ with finite support and any $\varphi \in H$ we have*

$$\sum_{m \in \ell} (Q^+\psi)_m \varphi_{n-m} = 0 \quad (33)$$

for all $n \in \ell$.

Proof. Denote by τ the following operator:

$$(\tau\psi)_n = \psi_{-n}.$$

We will have

$$\begin{aligned} \sum_{m \in \ell} (Q^+\psi)_m \varphi_{n-m} &= \sum_{m \in \ell} (Q^+\psi)_m (\tau\varphi)_{m-n} \\ &= \sum_{m \in \ell} \psi_m (Q\tau\varphi)_{m-n} \\ &= \sum_{m \in \ell} \psi_m (\tau Q^+\varphi)_{m-n} = 0. \end{aligned}$$

We used here the relation $\tau Q \tau = Q^+$ and the fact that the operators Q and Q^+ are formally conjugate to each other. $\square$

Corollary 3. *The assertion of Proposition 4 takes place for an arbitrary function G satisfying (28).*

Appendix 1

Proof of Theorem 1. Any representation $\rho : \pi_1(M) \rightarrow GL(2, \mathbb{R})$ can be obtained by specifying a flat linear connection in the trivial two-dimensional vector bundle over the 1-skeleton of $\mathcal{T}$. This means that, to each oriented edge $\langle P, P' \rangle$, we associate a 2×2 matrix $R_{P, P'}$ so that, for any triangle $\langle P, P', P'' \rangle$ of $\mathcal{T}$, we have $R_{P, P''} = R_{P, P'} R_{P', P''}$ and $R_{P, P'} = R_{P', P}^{-1}$. For a fixed vertex P_0 , a representation of $\pi_1(M, P_0)$ is obtained by putting $\rho(\gamma) = R_{P_0 P_1} \cdot \dots \cdot R_{P_{m-1} P_m} R_{P_m P_0}$, where the path γ consists of the edges $\langle P_0, P_1 \rangle, \dots, \langle P_{m-1}, P_m \rangle, \langle P_m, P_0 \rangle$. Clearly, any representation can be obtained in this way. Moreover, the representation is not changed under a “gauge transformation” $R_{P, P'} \mapsto C_P R_{P, P'} C_{P'}^{-1}$, where C_P , $P \neq P_0$, are arbitrary 2×2 matrices, and $C_{P_0} = \text{id}$.

Let us fix a vertex P'_0 connected with P_0 by an edge of $\mathcal{T}$. By a gauge transformation, we can achieve $R_{P_0 P'_0} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Let D be a simply connected domain containing P_0, P'_0 . Consider a local covariantly constant section of the vector bundle, *i.e.*, a vector function $v_P = (v_{P1}, v_{P2})$ on vertices from D such that $v_P \cdot R_{P, P'} = v_{P'}$ for any edge $\langle P, P' \rangle \subset D$. Since we have $v_{P_0 2} = v_{P'_0 1}$, the section v_P can be recovered from the knowledge of $v_{P_0 1}, v_{P'_0 1}$. The space of covariantly constant local sections is two-dimensional.

By forgetting the second coordinate, we obtain a two-dimensional space of functions v_{P1} , which can be turned into the space of covariant constants of a flat discrete connection associated with $\mathcal{T}$. This can be done by putting

$$b_{T, P_1} = c_{23}, \quad b_{T, P_2} = c_{31} d_{23}, \quad b_{T, P_3} = c_{12} d_{21}, \quad (34)$$

where $T = \langle P_1, P_2, P_3 \rangle$ is a triangle of $\mathcal{T}$ with a fixed enumeration of vertices, $c_{ij} = (R_{P_i, P_j})_{21}$, $d_{ij} = \det(R_{P_i, P_j})$. Changing the enumeration will result in the multiplication of all three coefficients (34) by the same constant, which has no effect on the corresponding discrete connection. Applying a generic gauge

transformation with $C_{P_0} = \text{id}$ we can always make all the coefficients (34) be non-zero.

The discrete connection $\{b_{T,P}\}$ obtained in this way has the following property. If we set $\psi_{P_0}, \psi_{P'_0}$ arbitrarily and extend ψ_P by solving equation (2) along a thick path, we will obtain the same function as if we set $v_{P_0} = (\psi_{P_0}, \psi_{P'_0})$, apply the parallel transport along the same path by using the connection R , and then forget the second coordinate of the obtained vector function. $\square$

Appendix 2

Here we exhibit a natural generalization of the constructions of Section 1.

Let X be a simplicial complex, $k \geq 1$ an integer. With any family $\mathcal{K}$ of k -dimensional simplices and numeric non-zero coefficients $b_{\sigma^k, P}$ defined for any simplex $\sigma^k \in \mathcal{K}$ and its vertex $P \in \sigma^k$ we associate the operator

$$(Q\psi)_{\sigma^k} = \sum_{P \in \sigma^k} b_{\sigma^k, P} \psi_P, \quad (35)$$

which takes functions of a vertex to functions on $\mathcal{K}$.

An analogue of discrete connections of Section 1 can be defined if $\mathcal{K}$ is the set of all k -simplices of X and X satisfies certain restrictions, which we now describe.

By a *k-thick path* in X we shall mean a sequence $\sigma_1, \sigma_2, \dots, \sigma_m$ of k -dimensional simplices such that $\sigma_i \cap \sigma_{i+1}$ is a $(k-1)$ -dimensional simplex for any $i = 1, \dots, m-1$. We call a thick path $\sigma_1, \dots, \sigma_m$ *elementary* if all simplices σ_i belong to the star of a single vertex. If a thick path γ_1 is obtained from a thick path γ_2 by an insertion or a removal an elementary path we say that the passage $\gamma_1 \mapsto \gamma_2$ is an *elementary homotopy*.

Any thick path $\sigma_1, \dots, \sigma_m$ defines a homotopy class of ordinary paths from the center of σ_1 to the center of σ_m in the obvious way. Thus, it makes sense to speak about homotopic thick paths.

We say that a simplicial complex X is *k-admissible* if any path is *k-thickable* and any homotopy between *k-thick* paths can be realized as a composition of elementary homotopies. For example, the *k-skeleton* of a triangulated manifold is always *k-admissible*.

For any *k-thick* loop $\gamma = (\sigma_1, \dots, \sigma_m, \sigma_{m+1} = \sigma_1)$ we define the *holonomy operator* R_γ in the same way as we did in Section 1. R_γ is a linear operator

on the space of solutions of the equation $Q\psi = 0$ in the simplex σ_1 . We say that the connection $\{b_{\sigma,P}\}$ has *trivial local holonomy* (or *zero curvature*) if $R_\gamma = \text{id}$ for any elementary k -thick loop γ .

Lemma 8. *If X is a k -admissible complex, then for any k -simplex σ_0^k the global holonomy of any discrete connection $\{b_{\sigma^k,P}\}$ with zero curvature along k -thick loops based at σ_0^k is a well defined homomorphism $\pi_1(X, \sigma_0^k) \rightarrow \mathbb{R}^k$.*

The proof is easy.

Consider the canonical connection associated with the family of all k -simplices of X , $b_{\sigma^k,P} \equiv 1$. It is easy to see that the canonical connection necessarily has trivial local holonomy at any vertex in the following two cases:

- 1) $k = 1$;
- 2) $k \geq 2$, X is a triangulated k -manifold such that any $(k-2)$ -dimensional simplex σ^{k-2} (*i.e.*, the number of k -simplices containing σ^{k-2}) is even.

We will assume one of these in the sequel. The operator $L = Q^+Q$ has the form

$$(L\psi)_P = \sum_{P'} m_{P,P'} \psi_{P'} + n_P \psi_P,$$

where the sum is taken over all vertices $P' \neq P$ from the star of P , $m_{P,P'}$ is the number of k -simplices containing the edge $\langle PP' \rangle$, n_P is the number of k -simplices containing the vertex P .

By analogy with Propositions 1, 2 one can prove the following.

Proposition 5. *The holonomy group $\mathcal{G}$ of the canonical connection of a triangulated k -manifold X all whose $(k-2)$ -simplices have even valence is a subgroup of the permutation group S_{k+1} .*

*Let q be the number of orbits of the action of $\mathcal{G} \subset S_{k+1}$ on the set $\{0, 1, \dots, k\}$. Then the dimension of the space of covariant constants of the canonical connections, *i.e.*, solutions of the equation $Q\psi = 0$ is $(q-1)$ -dimensional.*

The space of covariant constants coincides with the space of zero modes of the operator L .

In the particular case $k = 1$, zero modes of the operator L exist if and only if the 1-skeleton of X is a dichromatic graph. Now we suppose $k \geq 2$.

Consider the following three homomorphisms $\pi_1(X) \rightarrow \mathbb{Z}_2$:

- 1) the parity ρ_1 of the global holonomy of the canonical connection;
- 2) the orientation homomorphism ρ_2 ;
- 3) the parity homomorphism ρ_3 of the number of k -simplices in a k -thick path.

By analogy with Lemmas 2, 3, we have the following.

Lemma 9. *A triangulated k -manifold all whose $(k - 2)$ -simplices have even valence admits a b/w coloring of k -simplices such that any two adjacent k -simplices are of different color if and only if the representation ρ_3 is trivial.*

We have $\rho_3 = \rho_1\rho_2$.

If X admits a b/w coloring of k -simplices, then the operator L can also be factorized as follows:

$$L = 2Q_b^+Q_b = 2Q_w^+Q_w,$$

where Q_b, Q_w are operators associated with the family of black and white simplices, respectively. It is natural to ask whether the maximum principle holds for these operators. This question has not yet been investigated.

References

- [1] S. P. Novikov and I. A. Dynnikov, Discrete spectral symmetries of low-dimensional differential operators and difference operators on regular lattices and two-manifolds, (Russian) *Uspekhi Mat. Nauk* **52** (1997), no. 5, 175–234; translation in *Russian Math. Surveys* **52** (1997), no. 5, 1057–1116.
- [2] I. A. Dynnikov and S. P. Novikov, Laplace transformations and simplicial connections, (Russian) *Uspekhi Mat. Nauk* **52** (1997), no. 6, 157–158; translation in *Russian Math. Surveys* **52** (1997), no. 6, 1294–1295.
- [3] S. P. Novikov, Algebraic properties of two-dimensional difference operators, (Russian) *Uspekhi Mat. Nauk* **52** (1997), no. 1, 225–226; translation in *Russian Math. Surveys* **52** (1997), no. 1, 226–227
- [4] P. Sarnak, *Some applications of modular forms*, Camb. Univ. Press, 1990.